

Dawada neefta

(Somali)

Adiga iyo ilmahaagu waa in aad taqaanaan magaca daawo kasta oo ilmahaagu qaadanayo. Si gaar ah ayaa neeftu qofka walba u qabataa, daawada qof wax tartaana waxaa suurtoowda in aysan qofka kale wax tarin. Mar walba isticmaal tirada iyo qaabka daawooyinka laguugu qoray. Hubi in aad mar walba haysatid daawo kugu filan.

Asthma medicines

(English)

You and your child should know the name of each medicine your child is taking. Each person's asthma is different and medicines that help one person may not help another. Always use medicines in the amount and method prescribed. Always make sure you have enough medicines on hand.

Canaasiirta lagu neefsado oo deg dega u howlgasha (daawooyinka badbaadinta ama xanuunbi'inta)

Quick acting inhaled bronchodilators (rescue or reliever medicines)

Magac la yaqaan <i>Brand name</i>	Magac guud <i>Generic name</i>	Waxyaalaha loo baahanyahay in aad taqaan	Things to know	Waxyeelooyinka	Side effects
Maxair®	pirbuterol	<ul style="list-style-type: none"> Waxay furtaa tuubooyinka hawada iyadoo dabcinaysa muruqyada ku hareeraaran tuubooyinka hawada. 	<ul style="list-style-type: none"> Opens airways by relaxing the muscles around the airways. 	<ul style="list-style-type: none"> Gacmo jarceeya, “daganaasho la’aan” dareemaya, wadna garaac badan, madax xanuun. 	<ul style="list-style-type: none"> Shaky hands, “hyper” feeling, fast heartbeat, headache
Proventil® HFA Ventolin®	albuterol	<ul style="list-style-type: none"> Waa iyada in loogu hormariyaa haddii daawooyin kale oo neefta lala qaadanayo. 	<ul style="list-style-type: none"> Should be used first if other inhalers are taken at the same time. 	<ul style="list-style-type: none"> Waxyeelooyinku waxay jirayaan wakhti gaaban. Waxayna u badantahay in ay baaba’aan markii aad daawada si joogta ah aad u qaadatid. 	<ul style="list-style-type: none"> Side effects usually only last a short time. They will probably go away after using the medicine regularly.
Xopenex®	levalbuterol	<ul style="list-style-type: none"> Iskuday in aad u dhaxaysiisid 1 daqiiqo isku buufinta daawada, si ay midda ugu horaysaa u bilowdo shaqadeeda. Daawooyinkani waxay shaqadoodu bilaabataa 5 daqiiqo gudahood, waxayna jirka ku jirtaa 4 ilaa 6 saac. 	<ul style="list-style-type: none"> Try to wait 1 minute between puffs to let first puff begin working. These medicines start working in about 5 minutes and last 4 to 6 hours. 	<ul style="list-style-type: none"> Takhtarka wac haddii waxyeelooyinku dhib u hayaan ama baabiwaydo. 	<ul style="list-style-type: none"> Call doctor if the side effects are too bothersome or do not stop.

Canaasiirta lagu neefsado oo muddada dheer shaqaysa (ilaaliyaal)

Long-acting inhaled bronchodilators (controllers)

Magac la yaqaan <i>Brand name</i>	Magac guud <i>Generic name</i>	Waxyaalaha loo baahanyahay in aad taqaan	Things to know	Waxyeelooyinka	Side effects
Foradil®	formoterol	<ul style="list-style-type: none"> Waxay furtaa tuubooyinka hawada iyada oo dabcinaysa muruqyada ku heeraaran tuubooyinka hawada. Qaado 12kii saacba mar. Ha qaadan wax ka badan laba jeer maalintii. Marna ha u isticmaalin in ay bi'iso calaamada xanuunka Waa in aad ku neefsataa corticosteroid 	<ul style="list-style-type: none"> Opens airways by relaxing muscles around the airways. Take about 12 hours apart. Do not take more than twice a day. Never use to relieve symptoms. Should also be on an inhaled corticosteroid 	<ul style="list-style-type: none"> Waxay la midtahay midda kore. 	<ul style="list-style-type: none"> Same as above
Serevent®	salmeterol				

Naqaslaha istirowdku ku jiro (ilaaliyaal) *Inhaled corticosteroids (controllers)*

Magac la yaqaan <i>Brand name</i>	Magac guud <i>Generic name</i>	Waxyaalaha loo baahanyahay in aad taqaan	Things to know	Waxyeelooyinka	Side effects
Azmacort®	triamcinolone	<ul style="list-style-type: none"> Waxay ka hortagaysaa barar, jiro iyo xaako ku dhalata gudaha hawo mareenada. Wad isticmaalka daawadan maalin walba si aad uga hortagtid jirooyinka. Ha joojin qaadashada daawadan ilaa takhtar ku yiraahdo jooji. 	<ul style="list-style-type: none"> Prevents swelling, inflammation, and mucus inside the airways Continue using this medicine every day to prevent episodes. Do not stop taking this medicine until the doctor tells you to stop. 	<ul style="list-style-type: none"> Xabeeb Waxaa suurto gal ah cabeeb; ka luqluqo kadibna marka aad isticmaashid daawada, tuf si aad cudurka u ilaalisid in uusan kugu dhicin. Waa in aad la isticmaashaa isku-xiraha xiran (haddii aysan ahyn naqaslaha bool-bare la isticmaalo). 	<ul style="list-style-type: none"> Hoarseness Possible thrush; rinse mouth and spit after each use to prevent it. Must be used with a spacer (except if using a dry powder inhaler)
Azmanex®	mometasone				
Flovent®	fluticasone				
Pulmicort®	budesonide				
QVAR®	beclomethasone				

Lagu neefsada corticosteroid ku jiro/Cunsurka muddada dheer shaqaysa isku dhafanayaasha (ilaaliyaal)
Inhaled corticosteroid / Long-acting bronchodilator combinations (controllers)

Magac la yaqaan <i>Brand name</i>	Magac guud <i>Generic name</i>	Waxyaalaha loo baahanyahay in aad taqaan	Things to know	Waxyeelooyinka	Side effects
Advair® Symbicort®	fluticasone iyo salmeterol budesonide iyo formoterol	<ul style="list-style-type: none"> Waxay ka hortagaysaa barar, jiro iyo xaako ka dhalata gudaha hawo mareenada. Waxay furaysaa hawo mareenada iyaga oo dabcinaaya muruqyada ku heeraaran. Qaado maalin walba si aad uga hor tagtid cudurka. Ha joojin qaadashada daawadan ilaa takhtar ku yiraahdo moo'ye. Marna ha u isticmaalin in ay bi'iso calaamadaha cudurka. U dhaxaysii 12 saac daawada qaadashadeeda. Ha qaadan labo jeer ka badan maallintii. 	<ul style="list-style-type: none"> Prevents swelling, inflammation, and mucus inside the airways Opens airways by relaxing muscles around them Take every day to prevent episodes. Do not stop taking this medicine until the doctor tells you to stop. Never use it to relieve symptoms. Take doses about 12 hours apart. Do not take more than twice a day. 	<ul style="list-style-type: none"> Habeeb Waxaa suurto gal ah cabeeb, ka luquqo kadib marka aad isticmaashid daawada, tuf diyaha si aad cudurka u ilaalisid in uusan kugu dhicin. 	<ul style="list-style-type: none"> Hoarseness Possible thrush; rinse mouth and spit after each use to prevent it.

Naqaslaha leh daawada jiro ka hortagga (ilaaliyaal)

Inhaled anti-inflammatory medicines (controllers)

Magac la yaqaan <i>Brand name</i>	Magac guud <i>Generic name</i>	Waxyaalaha loo baahanyahay in aad taqaan	Things to know	Waxyeelooyinka	Side effects
Intal® Tilade®	cromolyn nedocromil	<ul style="list-style-type: none"> Waxay ka hortagaysaa jiro iyada oo isbadalka ka ilaalinaysa dahaarka tuubooyinka hawada, xiraysana qunbacasho dhacda. Qaado maalin walba si aad uga hortagtid jirada. Ha qaadan 4 jeer ka badan maallintii. Ha joojin qadashada daawadan ilaa takhtar ku yiraahdo jooji. 	<ul style="list-style-type: none"> Prevents inflammation by stabilizing cells that line the airway; blocks the reaction to triggers. Take every day to prevent episodes. Do not take more than 4 times daily. Do not stop taking this medicine until your doctor tells you to stop. 	<ul style="list-style-type: none"> Waxyeelooyinku waa naadir 	<ul style="list-style-type: none"> Side effects are very rare

Leukotriene qaabilaha caabiya (ilaaliye)		<i>Leukotriene receptor antagonist (controller)</i>			
Magac la yaqaan <i>Brand name</i>	Magac guud <i>Generic name</i>	Waxyaalaha loo baahanyahay in aad taqaan	Things to know	Waxyeeelooyinka	Side effects
Singulair®	montelukast	<ul style="list-style-type: none"> Waxay ka hotagaysaa barar, jirada, iyo xaakada. Qaado maalin walba si aad uga hortagtid jirada. Marna ha u isticmaalin in bi'iso calaamadaha cudurka. Ha joojin qaadashada daawadan ilaa takhtarku ku yiraahdo jooji. 	<ul style="list-style-type: none"> Prevents swelling, inflammation, and mucus. Take every day to prevent episodes. Never use to relieve symptoms. Do not stop taking this medicine until your doctor tells you to stop. 	<ul style="list-style-type: none"> Mahax xanuun ama callol xanuun Wac takhtarka haddii aad aragtid calaamadaha hargabka 	<ul style="list-style-type: none"> Headache or stomachache Call the doctor if flu-like symptoms occur.

Corticosteroids afka la qaato		<i>Oral corticosteroids</i>			
Magac la yaqaan <i>Brand name</i>	Magac guud <i>Generic name</i>	Waxyaalaha loo baahanyahay in aad taqaan	Things to know	Waxyeeelooyinka	Side effects
Orapred® Pediapred® Prelone® badeecooyin kala duwan <i>(various brands)</i>	prednisolone prednisone	<ul style="list-style-type: none"> Waxay yaraysaa bararka, jirada, iyo duufka ku jira tuubooyinka hawada. U qaado sida takhtarku kuugu qoray. Waxay shaqo biloydaa 6 ilaa 12 saac kadib. 	<ul style="list-style-type: none"> Decreases swelling, inflammation, and mucus inside the airways Take exactly as prescribed by your doctor. Starts to work in 6 to 12 hours 	<ul style="list-style-type: none"> Waxay furtaa cuntada, calool xanuun, dabciga oo isbadala, dareeraha oo ceshatid, waji dhalaala, iyo kala gowsyada oo xanuuna kadib marka aad joojisid qadashadeeda. Muddo dheer oo la qaataa waxay leedahay waxyeeelooyin kale, kala hadal dhakhtarkaaga. La qaado cunto ama caano si aad uga fogaatid calool xanuun 	<ul style="list-style-type: none"> Increased appetite, stomachache, mood changes, fluid retention, flushed face; perhaps temporary joint pain after it is stopped Long-term use may cause other side effects; talk to your doctor. Take with food or milk to avoid stomachache.

Su'aalo?

Warqadaani gaar uma aha ilmahaaga, laakiin waa warbixin guud. Haddii aad qabtid wax su'aal ah, fadlan wac bukaan socod eegtada (clinic) ama famashiyaha.

Questions?

This sheet is not specific to your child, but provides general information. If you have any questions, please call the clinic or pharmacy.

Children's Hospitals and Clinics of Minnesota
 Patient/Family Education
 2525 Chicago Avenue South
 Minneapolis, MN 55404
 5/07 ©Copyright, Translated 6/07